



# BRIGHT FUTURES HANDOUT ► PARENT

## 2 MONTH VISIT

Here are some suggestions from Bright Futures experts that may be of value to your family.

### ✓ HOW YOUR FAMILY IS DOING

- If you are worried about your living or food situation, talk with us. Community agencies and programs such as WIC and SNAP can also provide information and assistance.
- Find ways to spend time with your partner. Keep in touch with family and friends.
- Find safe, loving child care for your baby. You can ask us for help.
- Know that it is normal to feel sad about leaving your baby with a caregiver or putting him into child care.

### ✓ HOW YOU ARE FEELING

- Take care of yourself so you have the energy to care for your baby.
- Talk with me or call for help if you feel sad or very tired for more than a few days.
- Find small but safe ways for your other children to help with the baby, such as bringing you things you need or holding the baby's hand.
- Spend special time with each child reading, talking, and doing things together.

### ✓ FEEDING YOUR BABY

- Feed your baby only breast milk or iron-fortified formula until she is about 6 months old.
- Avoid feeding your baby solid foods, juice, and water until she is about 6 months old.
- Feed your baby when you see signs of hunger. Look for her to
  - Put her hand to her mouth.
  - Suck, root, and fuss.
- Stop feeding when you see signs your baby is full. You can tell when she
  - Turns away
  - Closes her mouth
  - Relaxes her arms and hands
- Burp your baby during natural feeding breaks.

#### If Breastfeeding

- Feed your baby on demand. Expect to breastfeed 8 to 12 times in 24 hours.
- Give your baby vitamin D drops (400 IU a day).
- Continue to take your prenatal vitamin with iron.
- Eat a healthy diet.
- Plan for pumping and storing breast milk. Let us know if you need help.
  - If you pump, be sure to store your milk properly so it stays safe for your baby. If you have questions, ask us.

#### If Formula Feeding

- Feed your baby on demand. Expect her to eat about 6 to 8 times each day, or 26 to 28 oz of formula per day.
- Make sure to prepare, heat, and store the formula safely. If you need help, ask us.
- Hold your baby so you can look at each other when you feed her.
- Always hold the bottle. Never prop it.

### ✓ YOUR GROWING BABY

- Have simple routines each day for bathing, feeding, sleeping, and playing.
- Hold, talk to, cuddle, read to, sing to, and play often with your baby. This helps you connect with and relate to your baby.
- Learn what your baby does and does not like.
- Develop a schedule for naps and bedtime. Put him to bed awake but drowsy so he learns to fall asleep on his own.
- Don't have a TV on in the background or use a TV or other digital media to calm your baby.
- Put your baby on his tummy for short periods of playtime. Don't leave him alone during tummy time or allow him to sleep on his tummy.
- Notice what helps calm your baby, such as a pacifier, his fingers, or his thumb. Stroking, talking, rocking, or going for walks may also work.
- *Never hit or shake your baby.*

### Helpful Resources:

Information About Car Safety Seats: [www.safercar.gov/parents](http://www.safercar.gov/parents) | Toll-free Auto Safety Hotline: 888-327-4236

## 2 MONTH VISIT—PARENT



### SAFETY

- Use a rear-facing-only car safety seat in the back seat of all vehicles.
- Never put your baby in the front seat of a vehicle that has a passenger airbag.
- Your baby's safety depends on you. Always wear your lap and shoulder seat belt. Never drive after drinking alcohol or using drugs. Never text or use a cell phone while driving.
- Always put your baby to sleep on her back in her own crib, not your bed.
  - Your baby should sleep in your room until she is at least 6 months old.
  - Make sure your baby's crib or sleep surface meets the most recent safety guidelines.
- If you choose to use a mesh playpen, get one made after February 28, 2013.
- Swaddling should not be used after 2 months of age.
- Prevent scalds or burns. Don't drink hot liquids while holding your baby.
- Prevent tap water burns. Set the water heater so the temperature at the faucet is at or below 120°F /49°C.
- Keep a hand on your baby when dressing or changing her on a changing table, couch, or bed.
- Never leave your baby alone in bathwater, even in a bath seat or ring.

## WHAT TO EXPECT AT YOUR BABY'S 4 MONTH VISIT

### We will talk about

- Caring for your baby, your family, and yourself
- Creating routines and spending time with your baby
- Keeping teeth healthy
- Feeding your baby
- Keeping your baby safe at home and in the car

Consistent with *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents*, 4th Edition

For more information, go to <https://brightfutures.aap.org>.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



The information contained in this handout should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances. Original handout included as part of the *Bright Futures Tool and Resource Kit*, 2nd Edition.

Inclusion in this handout does not imply an endorsement by the American Academy of Pediatrics (AAP). The AAP is not responsible for the content of the resources mentioned in this handout. Web site addresses are as current as possible but may change at any time.

The American Academy of Pediatrics (AAP) does not review or endorse any modifications made to this handout and in no event shall the AAP be liable for any such changes.

© 2019 American Academy of Pediatrics. All rights reserved.

# Your baby at 2 months



Baby's Name \_\_\_\_\_

Baby's Age \_\_\_\_\_

Today's Date \_\_\_\_\_

Milestones matter! How your baby plays, learns, speaks, acts, and moves offers important clues about his or her development. Check the milestones your baby has reached by 2 months. Take this with you and talk with your baby's doctor at every well-child visit about the milestones your baby has reached and what to expect next.

## What most babies do by this age:

### Social/Emotional Milestones

- Calms down when spoken to or picked up
- Looks at your face
- Seems happy to see you when you walk up to her
- Smiles when you talk to or smile at her

### Language/Communication Milestones

- Makes sounds other than crying
- Reacts to loud sounds

### Cognitive Milestones (learning, thinking, problem-solving)

- Watches you as you move
- Looks at a toy for several seconds

### Movement/Physical Development Milestones

- Holds head up when on tummy
- Moves both arms and both legs
- Opens hands briefly

## Other important things to share with the doctor...

- What are some things you and your baby do together?
- What are some things your baby likes to do?
- Is there anything your baby does or does not do that concerns you?
- Has your baby lost any skills he/she once had?
- Does your baby have any special healthcare needs or was he/she born prematurely?

**You know your baby best.** Don't wait. If your baby is not meeting one or more milestones, has lost skills he or she once had, or you have other concerns, act early. Talk with your baby's doctor, share your concerns, and ask about developmental screening. If you or the doctor are still concerned:

1. Ask for a referral to a specialist who can evaluate your baby more; and
2. Call your state or territory's early intervention program to find out if your baby can get services to help. Learn more and find the number at [cdc.gov/FindEI](https://www.cdc.gov/FindEI).

For more on how to help your baby, visit [cdc.gov/Concerned](https://www.cdc.gov/Concerned).

**Don't wait.  
Acting early can make  
a real difference!**



Download CDC's  
free Milestone  
Tracker app



American Academy  
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN®

# Help your baby learn and grow

As your baby's first teacher, you can help his or her learning and brain development. Try these simple tips and activities in a safe way. Talk with your baby's doctor and teachers if you have questions or for more ideas on how to help your baby's development.



- Respond positively to your baby. Act excited, smile, and talk to him when he makes sounds. This teaches him to take turns “talking” back and forth in conversation.
- Talk, read, and sing to your baby to help her develop and understand language.
- Spend time cuddling and holding your baby. This will help him feel safe and cared for. You will not spoil your baby by holding or responding to him.
- Being responsive to your baby helps him learn and grow. Limiting your screen time when you are with your baby helps you be responsive.
- Take care of yourself. Parenting can be hard work! It's easier to enjoy your new baby when you feel good yourself.
- Learn to notice and respond to your baby's signals to know what she's feeling and needs. You will feel good and your baby will feel safe and loved. For example, is she trying to “play” with you by making sounds and looking at you, or is she turning her head away, yawning, or becoming fussy because she needs a break?
- Lay your baby on his tummy when he is awake and put toys at eye level in front of him. This will help him practice lifting his head up. Do not leave your baby alone. If he seems sleepy, place him on his back in a safe sleep area (firm mattress with no blankets, pillows, bumper pads, or toys).
- Feed only breast milk or formula to your baby. Babies are not ready for other foods, water or other drinks for about the first 6 months of life.
- Learn when your baby is hungry by looking for signs. Watch for signs of hunger, such as putting hands to mouth, turning head toward breast/bottle, or smacking/licking lips.
- Look for signs your baby is full, such as closing her mouth or turning her head away from the breast/bottle. If your baby is not hungry, it's ok to stop feeding.
- Do not shake your baby or allow anyone else to—ever! You can damage his brain or even cause his death. Put your baby in a safe place and walk away if you're getting upset when he is crying. Check on him every 5–10 minutes. Infant crying is often worse in the first few months of life, but it gets better!
- Have routines for sleeping and feeding. This will help your baby begin to learn what to expect.

**To see more tips and activities download CDC's Milestone Tracker app.**

This milestone checklist is not a substitute for a standardized, validated developmental screening tool. These developmental milestones show what most children (75% or more) can do by each age. Subject matter experts selected these milestones based on available data and expert consensus.

[www.cdc.gov/ActEarly](http://www.cdc.gov/ActEarly) | 1-800-CDC-INFO (1-800-232-4636)



Download CDC's  
free Milestone  
Tracker app



**Learn the Signs. Act Early.**

# 2 months old

Child's name \_\_\_\_\_

Length \_\_\_\_\_ Weight \_\_\_\_\_ Date \_\_\_\_\_

Weight for Length percentile \_\_\_\_\_ %

## Food for Thought

*Any feeding concerns?*

*How is breastfeeding going?*

*Are you putting anything in your baby's bottle besides breast milk or formula?*

*How do you tell when your baby is hungry; when your baby is full?*

*Does your baby spend time on his or her belly?*

## Feeding Advice

- **Breastfeeding is highly recommended.**  
The best food for your baby is breastmilk.
- As your baby gets older and bigger, he or she will go longer between feedings and take more at each feeding.
- Expect to breastfeed your baby every 2-3 hours. If you are using formula, most babies take 4 oz. and feed 4-5 times per day. Total formula intake should be 16-24 ounces in 24 hours.
- Not all crying means hunger.
- Continue to use a 4 oz. bottle.
- Wait until at least 4-6 months to start cereal when your baby is developmentally ready.
- **DO NOT** put cereal in the bottle.
- **DO NOT** give your baby honey, juice, sweetened water or soft drinks.
- Ask about Vitamin D supplementation, especially for breastfed infants. For more information talk to your doctor or healthcare professional.


## Be Active

- Use crib mobiles and encourage reaching, kicking, stretching and belly play time.
- Limit time in swings and infant seats.
- Screen time (TV, computer, electronic games) not recommended under age 2.

## Notes:



The Ounce of Prevention Program is a collaboration of the Ohio Department of Health, Healthy Ohio; the American Academy of Pediatrics—Ohio Chapter; Nationwide Children's Hospital; the American Dairy Association Mideast and the Ohio Dietetic Association. May be reproduced in its entirety for educational purposes. February 2010





## BIRTH TO 6 MONTHS

### Safety for Your Child

Did you know that hundreds of children younger than 1 year die every year in the United States because of injuries—most of which could be prevented?

Often, injuries happen because parents are not aware of what their children can do. Children *learn quickly*, and before you know it, your child will be *wiggling* off a bed or *reaching* for your cup of hot coffee.

#### Car Injuries

**Car crashes** are a great threat to your child's life and health. Most injuries and deaths from car crashes **can be prevented** by the use of car safety seats. Your child, besides being much safer in a car safety seat, will behave better, so you can pay attention to your driving. Make your newborn's first ride home from the hospital a safe one—in a car safety seat. Your infant should ride in the back seat in a rear-facing car safety seat.

**Make certain that your baby's car safety seat is installed correctly. Read and follow the instructions that come with the car safety seat and the sections in the owners' manual of your car on using car safety seats correctly. Use the car safety seat EVERY time your child is in the car.**

NEVER put an infant in the front seat of a car with a passenger airbag.



#### Falls

Babies *wiggle* and *move* and *push* against things with their feet soon after they are born. Even these very first movements can result in a fall. As your baby grows and is able to roll over, he or she may fall off of things unless protected. **Do not leave your baby alone** on changing tables, beds, sofas, or chairs. **Put your baby in a safe place** such as a crib or playpen when you cannot hold him or her.

Your baby may be able to crawl as early as 6 months. **Use gates on stairways and close doors** to keep your baby out of rooms where he or she might get hurt. **Install operable window guards** on all windows above the first floor.

**Do not use a baby walker.** Your baby may tip the walker over, fall out of it, or fall down stairs and seriously injure his or her head. Baby walkers let children get to places where they can pull heavy objects or hot food on themselves. A better choice is a stationary activity center with no wheels.

**If your child has a serious fall or does not act normally after a fall, call your doctor.**



(over)

## Burns

At 3 to 5 months, babies will wave their fists and grab at things. **NEVER carry your baby and hot liquids, such as coffee, or foods at the same time.** You can't handle both. Your baby can get burned! To protect your child from tap water scalds, the hottest temperature at the faucet should be no more than 120°F. In many cases you can adjust your water heater.

**If your baby gets burned, immediately put the burned area in cold water. Keep the burned area in cold water for a few minutes to cool it off. Then cover the burn loosely with a dry bandage or clean cloth and call your doctor.**


To protect your baby from house fires, be sure you have a working smoke alarm on every level of your home, especially in furnace and sleeping areas. Test the alarms every month. It is best to use smoke alarms that use long-life batteries, but if you do not, change the batteries at least once a year.

## Choking and Suffocation

Babies explore their environment by putting anything and everything into their mouths. **NEVER leave small objects in your baby's reach, even for a moment.** NEVER feed your baby hard pieces of food such as chunks of raw carrots, apples, hot dogs, grapes, peanuts, and popcorn. Cut all the foods you feed your baby into thin pieces to prevent choking. **Be prepared if your baby starts to choke. Ask your doctor to recommend the steps you need to know. Learn how to save the life of a choking child.**

To prevent possible suffocation and reduce the risk of sudden infant death syndrome (SIDS), **your baby should always sleep on his or her back. Your baby should have his or her own crib or bassinet with no pillows, stuffed toys, bumpers, or loose bedding. NEVER put your baby on a waterbed, beanbag, or anything that is soft enough to cover the face and block air to the nose and mouth.**

**Plastic wrappers and bags** form a tight seal if placed over the mouth and nose and may suffocate your child. Keep them away from your baby.



From Your Doctor

The information in this publication should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances.

# Acetaminophen Dosing for Infants

This guide will help you use correct dosages, which ever product you have. This applies only to acetaminophen, and not to ibuprofen products.

Note: The products are packaged with different syringes, use the syringe that comes with the bottle.

## Infant + Children's Oral Suspension (160 mg/5mL)

Weight (lb)	Dose (mL)	Dose (tsp)
6-11 lbs	1.25 mL	¼ teaspoon
12-17 lbs	2.5 mL	½ teaspoon
18-23 lbs	3.75 mL	¾ teaspoon
24-35 lbs	5 mL	1 teaspoon



## SUGGESTED SCREEN TIME USE BY AGE



### 18 MONTHS AND YOUNGER

Avoid use of screen media other than video-chatting.



### 18 - 24 MONTHS

Parents of children 18 to 24 months of age who want to introduce digital media should choose high-quality programming, and watch it with their children to help them understand what they're seeing.



### 2 - 5 YEARS

Limit screen use to 1 hour per day of high-quality programs. Parents should co-view media with children to help them understand what they are seeing and apply it to the world around them.



### 6 - 12 YEARS

Place consistent limits on the time spent using media, and the types of media, and make sure media does not take the place of adequate sleep, physical activity and other behaviors essential to health.



### 12 YEARS AND OLDER

Designate media-free times together, such as dinner or driving, as well as media-free locations at home, such as bedrooms.

# Pneumococcal Conjugate Vaccine: *What You Need to Know*

Many vaccine information statements are available in Spanish and other languages. See [www.immunize.org/vis](http://www.immunize.org/vis)

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite [www.immunize.org/vis](http://www.immunize.org/vis)

## 1. Why get vaccinated?

Pneumococcal conjugate vaccine can prevent pneumococcal disease.

**Pneumococcal disease** refers to any illness caused by pneumococcal bacteria. These bacteria can cause many types of illnesses, including pneumonia, which is an infection of the lungs. Pneumococcal bacteria are one of the most common causes of pneumonia.

Besides pneumonia, pneumococcal bacteria can also cause:

- Ear infections
- Sinus infections
- Meningitis (infection of the tissue covering the brain and spinal cord)
- Bacteremia (infection of the blood)

Anyone can get pneumococcal disease, but children under 2 years old, people with certain medical conditions or other risk factors, and adults 65 years or older are at the highest risk.

Most pneumococcal infections are mild. However, some can result in long-term problems, such as brain damage or hearing loss. Meningitis, bacteremia, and pneumonia caused by pneumococcal disease can be fatal.

## 2. Pneumococcal conjugate vaccine

Pneumococcal conjugate vaccine helps protect against bacteria that cause pneumococcal disease. There are three pneumococcal conjugate vaccines (PCV13, PCV15, and PCV20). The different vaccines are recommended for different people based on age and medical status. Your health care provider can help you determine which type of pneumococcal conjugate vaccine, and how many doses, you should receive.

**Infants and young children** usually need 4 doses of pneumococcal conjugate vaccine. These doses are recommended at 2, 4, 6, and 12–15 months of age.

**Older children and adolescents** might need pneumococcal conjugate vaccine depending on their age and medical conditions or other risk factors if they did not receive the recommended doses as infants or young children.

**Adults 19 through 64 years old** with certain medical conditions or other risk factors who have not already received pneumococcal conjugate vaccine should receive pneumococcal conjugate vaccine.

**Adults 65 years or older** who have not previously received pneumococcal conjugate vaccine should receive pneumococcal conjugate vaccine.

Some people with certain medical conditions are also recommended to receive pneumococcal polysaccharide vaccine (a different type of pneumococcal vaccine, known as PPSV23). Some adults who have previously received a pneumococcal conjugate vaccine may be recommended to receive another pneumococcal conjugate vaccine.



---

### 3. Talk with your health care provider

---

Tell your vaccination provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of any type of pneumococcal conjugate vaccine (PCV13, PCV15, PCV20, or an earlier pneumococcal conjugate vaccine known as PCV7), or to any vaccine containing diphtheria toxoid (for example, DTaP), or has any severe, life-threatening allergies**

In some cases, your health care provider may decide to postpone pneumococcal conjugate vaccination until a future visit.

People with minor illnesses, such as a cold, may be vaccinated. People who are moderately or severely ill should usually wait until they recover.

Your health care provider can give you more information.

---

### 4. Risks of a vaccine reaction

---

- Redness, swelling, pain, or tenderness where the shot is given, and fever, loss of appetite, fussiness (irritability), feeling tired, headache, muscle aches, joint pain, and chills can happen after pneumococcal conjugate vaccination.

Young children may be at increased risk for seizures caused by fever after a pneumococcal conjugate vaccine if it is administered at the same time as inactivated influenza vaccine. Ask your health care provider for more information.

People sometimes faint after medical procedures, including vaccination. Tell your provider if you feel dizzy or have vision changes or ringing in the ears.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

---

### 5. What if there is a serious problem?

---

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at [www.vaers.hhs.gov](http://www.vaers.hhs.gov) or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff members do not give medical advice.*

---

### 6. The National Vaccine Injury Compensation Program

---

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Claims regarding alleged injury or death due to vaccination have a time limit for filing, which may be as short as two years. Visit the VICP website at [www.hrsa.gov/vaccinecompensation](http://www.hrsa.gov/vaccinecompensation) or call **1-800-338-2382** to learn about the program and about filing a claim.

---

### 7. How can I learn more?

---

- Ask your health care provider.
- Call your local or state health department.
- Visit the website of the Food and Drug Administration (FDA) for vaccine package inserts and additional information at [www.fda.gov/vaccines-blood-biologics/vaccines](http://www.fda.gov/vaccines-blood-biologics/vaccines).
- Contact the Centers for Disease Control and Prevention (CDC):
  - Call **1-800-232-4636 (1-800-CDC-INFO)** or
  - Visit CDC's website at [www.cdc.gov/vaccines](http://www.cdc.gov/vaccines).



# Your Child's First Vaccines: What You Need to Know

Many vaccine information statements are available in Spanish and other languages. See [www.immunize.org/vis](http://www.immunize.org/vis)

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite [www.immunize.org/vis](http://www.immunize.org/vis)

The vaccines included on this statement are likely to be given at the same time during infancy and early childhood. There are separate *Vaccine Information Statements* for other vaccines that are also routinely recommended for young children (measles, mumps, rubella, varicella, rotavirus, influenza, and hepatitis A)

**Your child is getting these vaccines today:**

- DTaP       Hib       Hepatitis B       PCV       Polio

(Provider: Check appropriate boxes.)

## 1. Why get vaccinated?

Vaccines can prevent disease. Childhood vaccination is essential because it helps provide immunity before children are exposed to potentially life-threatening diseases.

### Diphtheria, tetanus, and pertussis (DTaP)

- **Diphtheria (D)** can lead to difficulty breathing, heart failure, paralysis, or death.
- **Tetanus (T)** causes painful stiffening of the muscles. Tetanus can lead to serious health problems, including being unable to open the mouth, having trouble swallowing and breathing, or death.
- **Pertussis (aP)**, also known as “whooping cough,” can cause uncontrollable, violent coughing that makes it hard to breathe, eat, or drink. Pertussis can be extremely serious, especially in babies and young children, causing pneumonia, convulsions, brain damage, or death.

### Hib (*Haemophilus influenzae* type b) disease

*Haemophilus influenzae* type b can cause many different kinds of infections. Hib bacteria can cause mild illness, such as ear infections or bronchitis, or they can cause severe illness, such as infections of the blood. Hib infection can also cause pneumonia; severe swelling in the throat, making it hard to breathe; and infections of the blood, joints, bones, and covering of the heart. Severe Hib infection, also called “invasive Hib disease,” requires treatment in a hospital and can sometimes result in death.

### Hepatitis B

**Hepatitis B** is a liver disease that can cause mild illness lasting a few weeks, or it can lead to a serious, lifelong illness. Acute hepatitis B infection is a short-term illness that can lead to fever, fatigue, loss of appetite, nausea, vomiting, jaundice (yellow skin or eyes, dark urine, clay-colored bowel movements), and pain in the muscles, joints, and stomach. Chronic hepatitis B infection is a long-term illness that occurs when the hepatitis B virus remains in a person’s body. Most people who go on to develop chronic hepatitis B do not have symptoms, but it is still very serious and can lead to liver damage (cirrhosis), liver cancer, and death.

### Pneumococcal disease (PCV)

**Pneumococcal disease** refers to any illness caused by pneumococcal bacteria. These bacteria can cause many types of illnesses, including pneumonia, which is an infection of the lungs. Besides pneumonia, pneumococcal bacteria can also cause ear infections, sinus infections, meningitis (infection of the tissue covering the brain and spinal cord), and bacteremia (infection of the blood). Most pneumococcal infections are mild. However, some can result in long-term problems, such as brain damage or hearing loss. Meningitis, bacteremia, and pneumonia caused by pneumococcal disease can be fatal.



**U.S. Department of Health and Human Services**  
Centers for Disease Control and Prevention

## Polio

**Polio** (or poliomyelitis) is a disabling and life-threatening disease caused by poliovirus, which can infect a person's spinal cord, leading to paralysis. Most people infected with poliovirus have no symptoms, and many recover without complications. Some people infected with poliovirus will experience sore throat, fever, tiredness, nausea, headache, or stomach pain, and most people with these symptoms will also recover without complications. A smaller group of people will develop more serious symptoms: paresthesia (feeling of pins and needles in the legs), meningitis (infection of the covering of the spinal cord and/or brain), or paralysis (can't move parts of the body) or weakness in the arms, legs, or both. Paralysis can lead to permanent disability and death.

---

## 2. DTaP, Hib, hepatitis B, pneumococcal conjugate, and polio vaccines

---

**Infants and children** usually need:

- 5 doses of **diphtheria, tetanus, and acellular pertussis vaccine (DTaP)**
- 3 or 4 doses of **Hib vaccine**
- 3 doses of **hepatitis B vaccine**
- 4 doses of **pneumococcal conjugate vaccine (PCV)**
- 4 doses of **polio vaccine**

Some children might need fewer or more than the usual number of doses of some vaccines to have the best protection because of their age at vaccination or other circumstances.

**Older children, adolescents, and adults** with certain health conditions or other risk factors or who did not get vaccinated earlier might also be recommended to receive 1 or more doses of some of these vaccines.

These vaccines are given as either stand-alone vaccines or as part of a combination vaccine (a type of vaccine that combines more than one vaccine together into one shot).

---

## 3. Talk with your health care provider

---

Tell your vaccination provider if the child getting the vaccine:

### **For all of these vaccines:**

- Has had an **allergic reaction after a previous dose of the vaccine**, or has any **severe, life-threatening allergies**

### **For DTaP:**

- Has had an **allergic reaction after a previous dose of any vaccine that protects against diphtheria, tetanus, or pertussis**
- Has had a **coma, decreased level of consciousness, or prolonged seizures within 7 days after a previous dose of any pertussis vaccine (DTP or DTaP)**
- Has **seizures or another nervous system problem**
- Has ever had **Guillain-Barré syndrome** (also called "GBS")
- Has had **severe pain or swelling after a previous dose of any vaccine that protects against diphtheria or tetanus**

### **For PCV:**

- Has had an **allergic reaction after a previous dose of any type of pneumococcal conjugate vaccine (PCV13, PCV15, PCV20, or an earlier pneumococcal conjugate vaccine known as PCV7), or to any vaccine containing diphtheria toxoid** (for example, DTaP)

In some cases, your child's health care provider may decide to postpone vaccination until a future visit.

Children with minor illnesses, such as a cold, may be vaccinated. Children who are moderately or severely ill should usually wait until they recover before being vaccinated.

Your child's health care provider can give you more information.

---

## 4. Risks of a vaccine reaction

---

### For all of these vaccines:

- Soreness, redness, swelling, warmth, pain, or tenderness where the shot is given can happen after vaccination.

### For DTaP vaccine, Hib vaccine, hepatitis B vaccine, and PCV:

- Fever can happen after vaccination.

### For DTaP vaccine:

- Fussiness, feeling tired, loss of appetite, and vomiting sometimes happen after DTaP vaccination.
- More serious reactions, such as seizures, non-stop crying for 3 hours or more, or high fever (over 105°F) after DTaP vaccination happen much less often. Rarely, vaccination is followed by swelling of the entire arm or leg, especially in older children when they receive their fourth or fifth dose.

### For PCV:

- Loss of appetite, fussiness (irritability), feeling tired, headache, and chills can happen after PCV vaccination.
- Young children may be at increased risk for seizures caused by fever after a pneumococcal conjugate vaccine if it is administered at the same time as inactivated influenza vaccine. Ask your health care provider for more information.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

---

## 5. What if there is a serious problem?

---

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at [www.vaers.hhs.gov](http://www.vaers.hhs.gov) or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff members do not give medical advice.*

---

## 6. The National Vaccine Injury Compensation Program

---

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Claims regarding alleged injury or death due to vaccination have a time limit for filing, which may be as short as two years. Visit the VICP website at [www.hrsa.gov/vaccinecompensation](http://www.hrsa.gov/vaccinecompensation) or call **1-800-338-2382** to learn about the program and about filing a claim.

---

## 7. How can I learn more?

---

- Ask your health care provider.
- Call your local or state health department.
- Visit the website of the Food and Drug Administration (FDA) for vaccine package inserts and additional information at [www.fda.gov/vaccines-blood-biologics/vaccines](http://www.fda.gov/vaccines-blood-biologics/vaccines).
- Contact the Centers for Disease Control and Prevention (CDC):
  - Call **1-800-232-4636 (1-800-CDC-INFO)** or
  - Visit CDC's website at [www.cdc.gov/vaccines](http://www.cdc.gov/vaccines).



# Rotavirus Vaccine:

## *What You Need to Know*

Many vaccine information statements are available in Spanish and other languages. See [www.immunize.org/vis](http://www.immunize.org/vis)

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite [www.immunize.org/vis](http://www.immunize.org/vis)

### 1. Why get vaccinated?

Rotavirus vaccine can prevent rotavirus disease.

Rotavirus commonly causes severe, watery diarrhea, mostly in babies and young children. Vomiting and fever are also common in babies with rotavirus. Children may become dehydrated and need to be hospitalized and can even die.

### 2. Rotavirus vaccine

Rotavirus vaccine is administered by putting drops in the child's mouth. Babies should get 2 or 3 doses of rotavirus vaccine, depending on the brand of vaccine used.

- The first dose must be administered before 15 weeks of age.
- The last dose must be administered by 8 months of age.

Almost all babies who get rotavirus vaccine will be protected from severe rotavirus diarrhea.

Another virus called "porcine circovirus" can be found in one brand of rotavirus vaccine (Rotarix). This virus does not infect people, and there is no known safety risk.

Rotavirus vaccine may be given at the same time as other vaccines.

### 3. Talk with your health care provider

Tell your vaccination provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of rotavirus vaccine**, or has any **severe, life-threatening allergies**
- Has a **weakened immune system**
- Has **severe combined immunodeficiency (SCID)**
- Has had a type of bowel blockage called "**intussusception**"

In some cases, your child's health care provider may decide to postpone rotavirus vaccination until a future visit.

Infants with minor illnesses, such as a cold, may be vaccinated. Infants who are moderately or severely ill should usually wait until they recover before getting rotavirus vaccine.

Your child's health care provider can give you more information.

### 4. Risks of a vaccine reaction

- Irritability or mild, temporary diarrhea or vomiting can happen after rotavirus vaccine.

Intussusception is a type of bowel blockage that is treated in a hospital and could require surgery. It happens naturally in some infants every year in the United States, and usually there is no known reason for it. There is also a small risk of intussusception from rotavirus vaccination, usually within a week after the first or second vaccine dose. This additional risk is estimated to range from about 1 in 20,000 U.S. infants to 1 in 100,000 U.S. infants who get rotavirus vaccine. Your health care provider can give you more information.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.



**U.S. Department of Health and Human Services**  
Centers for Disease Control and Prevention

---

## 5. What if there is a serious problem?

---

For intussusception, look for signs of stomach pain along with severe crying. Early on, these episodes could last just a few minutes and come and go several times in an hour. Babies might pull their legs up to their chest. Your baby might also vomit several times or have blood in the stool, or could appear weak or very irritable. These signs would usually happen during the first week after the first or second dose of rotavirus vaccine, but look for them any time after vaccination. If you think your baby has intussusception, contact a health care provider right away. If you can't reach your health care provider, take your baby to a hospital. Tell them when your baby got rotavirus vaccine.

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at [www.vaers.hhs.gov](http://www.vaers.hhs.gov) or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff members do not give medical advice.*

---

## 6. The National Vaccine Injury Compensation Program

---

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Claims regarding alleged injury or death due to vaccination have a time limit for filing, which may be as short as two years. Visit the VICP website at [www.hrsa.gov/vaccinecompensation](http://www.hrsa.gov/vaccinecompensation) or call **1-800-338-2382** to learn about the program and about filing a claim.

---

## 7. How can I learn more?

---

- Ask your health care provider.
- Call your local or state health department.
- Visit the website of the Food and Drug Administration (FDA) for vaccine package inserts and additional information at [www.fda.gov/vaccines-blood-biologics/vaccines](http://www.fda.gov/vaccines-blood-biologics/vaccines).
- Contact the Centers for Disease Control and Prevention (CDC):
  - Call **1-800-232-4636** (**1-800-CDC-INFO**) or
  - Visit CDC's website at [www.cdc.gov/vaccines](http://www.cdc.gov/vaccines).





***Pregnant and new moms often  
feel anxious or depressed***

*perinatal outreach  
& encouragement*

**poem**  
*for  
moms*

A Program of Mental Health America of Ohio

***A free maternal mental health support  
program in Ohio***

**We offer pregnant and parenting women**

- a dedicated peer support and care navigation line
- mom-to-mom mentoring
- online and in-person support groups
- referrals to specialized maternal mental health clinicians
- peer support and referral services by and for African-American women
- referrals to support services for pregnancy and infant loss

***How to get help:***

***Call or text: 614.315.8989  
or visit: [poemonline.org](http://poemonline.org)***

## **Anxiety and depression are the most common complications of pregnancy and childbirth.**

Maternal mental health complications can appear during pregnancy or for many months after childbirth. These symptoms—from mild to severe—do not typically resolve without appropriate support and treatment.

“Postpartum depression” is often the term used to reference these symptoms, though they often do not appear as depression. Moms may have anxiety, panic, scary thoughts, anger, and mania, and all of these symptoms are on the spectrum of maternal mental health complications.

### **You might feel:**

Overwhelmed

Guilty

Anxious or nervous

Hopeless

Irritable or angry

Scared by your thoughts

Disconnected

Exhausted but unable to sleep

***POEM can help.***

**Call or text: 614.315.8989  
or visit: [poemonline.org](http://poemonline.org)**



*POEM is a program of the non-profit organization Mental Health America of Ohio and the Ohio chapter of Postpartum Support International, the leading authority on pregnancy & postpartum mental health.*

FREE BOOKS EACH  
**MONTH**

CHILDREN AGES  
**0-5**

REGISTER YOUR CHILD  
**TODAY**

# FREE BOOKS



**RECEIVE A FREE  
HIGH-QUALITY, AGE APPROPRIATE BOOK EACH  
MONTH FOR REGISTERED CHILDREN FROM  
BIRTH TO AGE FIVE.**

**REGISTER YOUR CHILD TODAY!  
OHIOIMAGINATIONLIBRARY.ORG/ENROLL**



**DOLLY PARTON'S IMAGINATION LIBRARY OF OHIO**

